



ACTIVE INGREDIENT

Imidacloprid 30.5% SC

RECOMMENDED USES

Can be used for management of termites for pre-construction as well as post-construction anti-termite treatment

MODE OF ACTION

Systemic insecticide with contact and stomach action

BENEFITS

- Non-repellent insecticide, hence termites fail to detect gaps in the chemical barrier and are directly exposed to the chemical even in improper application situation
- Premise dose not kill termites immediately on contact and allows more termites to get exposed to treated soil, thus giving a better control
- Exposed termited transfer toxicant to nest mates not directly exposed to termiticide, causing indirect mortality
- Odourless So no undesirable smell

DOSAGE

Dilute 2.1 ml of formulation in one litre of water for the control of termites in buildings during pre and post - construction anti-termite treatment

Note: Treatment has to be carried out as per the current BIS (IS 6313) recommendations

ANTIDOTE

There is no known specific antidote. Follow directions given in the pack leaflet. Treat symptomatically

PACKS

250 ml and 5 Litre



Insecticide for effective termite management



Corporate Office:

Crop Science - Environmental ScienceBayer House, Central Avenue, Hiranandani Estate
Thane (West) - 4000 607. Maharashtra, India.

Customer Care Cell: E-mail: customercare@bayer.com Tel.:022 - 2531 1234

Toll Free: 1800 209 5666

Choose Safety Choose Bayer



